

Name of the Course : CBCS B.Sc. Mathematical Sciences / B.Sc. (Prog.)
Unique Paper Code : 42357618
Name of the Paper : DSE- Numerical Methods
Semester : VI
Duration : 3 Hours
Maximum Marks : 75 Marks

Attempt any four questions. All questions carry equal marks. All symbols have usual meanings.

1. Let $f(x) = x^4 - 18x^2 + 45$ and $g(x) = x^3 + x^2 - 3x - 3$.
 - (i) Verify that both the equations $f(x) = 0$ and $g(x) = 0$ have a root on the interval (1, 2).
 - (ii) By performing three iterations of Newton-Raphson method, with $x_0 = 1$, find an approximation of the root of $f(x) = 0$.
 - (iii) By performing three iterations of Bisection method, find an approximation of the root of $g(x) = 0$.
 - (iv) Given that the exact value of the root in both cases is $x = \sqrt{3}$, compute the absolute error in the approximations obtained.

2. Find the inverse of A, using Gauss-Jordan elimination

$$A = \begin{bmatrix} 1 & 3 & 5 \\ 2 & 7 & 13 \\ 3 & 11 & 22 \end{bmatrix}.$$

Using Gauss-Seidel iteration method, solve the system of equations given by

$$A \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1.1 \\ 1.2 \\ 1.3 \end{bmatrix}.$$

3. Use the Lagrange's interpolation to find a polynomial that passes through the points (0, 2), (1, 3), (2, 12) and (5, 147).

Using the following data

| | | | | |
|--------|--------|--------|--------|---------|
| x | 1.0 | 1.5 | 2.0 | 2.5 |
| $f(x)$ | 2.7183 | 4.4817 | 7.3891 | 12.1825 |

estimate the value of $f(2.15)$ using

- (i) Newton's forward difference interpolation
- (ii) Newton's backward difference interpolation.

Compare the errors and find which of the methods gave a better approximation of $f(2.15)$.

4. The following data gives the velocity of a particle for 20 seconds at an interval of 5 seconds. Find the initial acceleration using the entire data:

| | | | | | |
|----------------------|---|---|----|----|-----|
| Time t (sec) | 0 | 5 | 10 | 15 | 20 |
| Velocity v (m/sec) | 0 | 2 | 13 | 68 | 227 |

A boundary value problem is defined by

$$\frac{d^2y}{dx^2} + y + 1 = 0, \quad 0 \leq x \leq 1$$

where $y(0) = 0$ and $y(1) = 0$ with $h = 0.5$. Use the finite difference method to determine the value of $y(0.5)$. Its exact solution is

$$y(x) = \cos x + \frac{1 - \cos 1}{\sin 1} \sin x - 1.$$

Calculate the error.

5. Use the formula

$$f'(x_0) \approx \frac{f(x_0) - f(x_0 - h)}{h}$$

to approximate the derivative of $f(x) = \sin x$ at $x_0 = \pi$ taking $h = 1, 0.1, 0.01$.

Use Euler's method to approximate the solution of the initial value problem

$$\frac{dy}{dt} = \frac{(1+y)^2}{t}, \quad y(1) = 0, 1 \leq t \leq 4$$

taking 5 steps.

6. Find approximate value of the integral $I = \int_0^2 e^x dx$ using
- Trapezoidal Rule;
 - Simpson's 1/3 rule;
 - Simpson's 3/8 rule.